

Literatur und Anmerkungen zum Beitrag von Robyn Cody, Jan-Niklas Kreppke, Oliver Faude und Markus Gerber „Was bewegt Depressive – Sport und Depression“, Kerbe 3/2020, Seite 37

- Holsboer-Trachsler E HJ, Beck J, Brand S, Hemmeter UM, Keck ME, et al. Die Akutbehandlung depressiver Episoden. *Swiss Medical Forum*. 2016;16:716–24.
- Stubbs B, Vancampfort D, Hallgren M, Firth J, Veronese N, Solmi M, et al. EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and Position Statement from the European Psychiatric Association (EPA), supported by the International Organization of Physical Therapists in Mental Health (IOPTMH). *Eur Psychiatry*. 2018;54:124–44.
- Knapen J, Vancampfort D, Morien Y, Marchal Y. Exercise therapy improves both mental and physical health in patients with major depression. *Disabil Rehabil*. 2015;37(16):1490–5.
- Stubbs B, Rosenbaum S, Vancampfort D, Ward PB, Schuch FB. Exercise improves cardiorespiratory fitness in people with depression: A meta-analysis of randomized control trials. *Journal of affective disorders*. 2016;190:249–53.
- Blumenthal J, Smith P, Hofmann B. Is exercise a viable treatment for depression? . *ACSMs Health Fit J*. 2012(16):14–21.
- Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S, Khatri P, et al. Effects of exercise training on older patients with major depression. *Arch Intern Med*. 1999;159(19):2349–56.
- Dimeo F, Bauer M, Varahram I, Proest G, Halter U. Benefits from aerobic exercise in patients with major depression: a pilot study. *British journal of sports medicine*. 2001;35(2):114–7.
- Singh NA, Clements KM, Singh MA. The efficacy of exercise as a long-term antidepressant in elderly subjects: a randomized, controlled trial. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2001;56(8):M497–504.
- Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomised controlled trials. *Bmj*. 2001;322(7289):763–7.
- Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA, et al. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosomatic medicine*. 2007;69(7):587–96.
- Cooney GM, Dwan K, Greig CA, Lawlor DA, Rimer J, Waugh FR, et al. Exercise for depression. *The Cochrane database of systematic reviews*. 2013(9):Cd004366.
- Mather AS, Rodríguez C, Guthrie MF, McHarg AM, Reid IC, McMurdo ME. Effects of exercise on depressive symptoms in older adults with poorly responsive depressive disorder: randomised controlled trial. *The British journal of psychiatry : the journal of mental science*. 2002;180:411–5.
- Schuch FB, Vasconcelos-Moreno MP, Borowsky C, Zimmermann AB, Rocha NS, Fleck MP. Exercise and severe major depression: effect on symptom severity and quality of life at discharge in an inpatient cohort. *Journal of psychiatric research*. 2015;61:25–32.
- Kvam S, Kleppe CL, Nordhus IH, Hovland A. Exercise as a treatment for depression: A meta-analysis. *Journal of affective disorders*. 2016;202:67–86.
- Rethorst CD, Wipfli BM, Landers DM. The antidepressive effects of exercise: a meta-analysis of randomized trials. *Sports medicine (Auckland, NZ)*. 2009;39(6):491–511.
- Mota-Pereira J, Silverio J, Carvalho S, Ribeiro JC, Fonte D, Ramos J. Moderate exercise improves depression parameters in treatment-resistant patients with major depressive disorder. *Journal of psychiatric research*. 2011;45(8):1005–11.
- Babyak M, Blumenthal JA, Herman S, Khatri P, Doraiswamy M, Moore K, et al. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosomatic medicine*. 2000;62(5):633–8.
- Hoffman BM, Babyak MA, Craighead WE, Sherwood A, Doraiswamy PM, Coons MJ, et al. Exercise and pharmacotherapy in patients with major depression: one-year follow-up of the SMILE study. *Psychosomatic medicine*. 2011;73(2):127–33.
- Mammen G, Faulkner G. Physical activity and the prevention of depression: a systematic review of prospective studies. *American journal of preventive medicine*. 2013;45(5):649–57.
- Callaghan P, Khalil E, Morres I, Carter T. Pragmatic randomised controlled trial of preferred intensity exercise in women living with depression. *BMC Public Health*. 2011;11(1):465.
- Helgadottir B, Hallgren M, Ekblom O, Forsell Y. Training fast or slow? Exercise for depression: A randomized controlled trial. *Preventive medicine*. 2016;91:123–31.
- Gerber M, Minghetti A, Beck J, Zahner L, Donath L. Sprint Interval Training and Continuous Aerobic Exercise Training Have Similar Effects on Exercise Motivation and Affective Responses to Exercise in Patients With Major Depressive Disorders: A Randomized Controlled Trial. *Front Psychiatry*. 2018;9:694–.
- Minghetti A, Faude O, Hanssen H, Zahner L, Gerber M, Donath L. Sprint interval training (SIT) substantially reduces depressive symptoms in major depressive disorder (MDD): A randomized controlled trial. *Psychiatry Research*. 2018;265:292–7.
- Nebiker L, Lichtenstein E, Minghetti A, Zahner L, Gerber M, Faude O, et al. Moderating Effects of Exercise Duration and Intensity in Neuromuscular vs. Endurance Exercise Interventions for the Treatment of Depression: A Meta-Analytical Review. *Front Psychiatry*. 2018;9:305.
- Gerber M, Minghetti A, Beck J, Zahner L, Donath L. Is improved fitness following a 12-week exercise program associated with decreased symptom severity, better wellbeing, and fewer sleep complaints in patients with major depressive disorders? A secondary analysis of a randomized controlled trial. *Journal of psychiatric research*. 2019;113:58–64.
- Schuch FB, Vancampfort D, Richards J, Rosenbaum S, Ward PB, Stubbs B. Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *Journal of psychiatric research*. 2016;77:42–51.
- Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *Jama*. 2009;301(19):2024–35.
- Correll CU, Solmi M, Veronese N, Bortolato B, Rossion S, Santonastaso P, et al. Prevalence, incidence and mortality from cardiovascular disease in patients with pooled and specific severe mental illness: a large-scale meta-analysis of 3,211,768 patients and 113,383,368 controls. *World psychiatry : official journal of the World Psychiatric Association (WPA)*. 2017;16(2):163–80.
- Lawrence D, Hancock KJ, Kisely S. The gap in life expectancy from preventable physical illness in psychiatric patients in Western Australia: retrospective analysis of population based registers. *BMJ : British Medical Journal*. 2013;346:f2539.
- Gerber M, Lindwall M, Lindegard A, Borjesson M, Jonsdottir IH. Cardiorespiratory fitness protects against stress-related symptoms of burnout and depression. *Patient education and counseling*. 2013;93(1):146–52.
- Helmich I, Latini A, Sigwalt A, Carta MG, Machado S, Velasques B, et al. Neurobiological alterations induced by exercise and their impact on depressive disorders [corrected]. *Clinical practice and epidemiology in mental health : CP & EMH*. 2010;6:115–25.
- Ludyga S. Sportaktivität, Stress und das Gehirn. In: Fuchs R, Gerber M, editors. *Stressregulation und Sport*. Heidelberg: Springer; 2017.
- Hamer M, Steptoe A. Association between physical fitness, parasympathetic control, and proinflammatory responses to mental stress. *Psychosomatic medicine*. 2007;69(7):660–6.
- Mucke M, Ludyga S, Colledge F, Gerber M. Influence of Regular Physical Activity and Fitness on Stress Reactivity as Measured with the Trier Social Stress Test Protocol: A Systematic Review. *Sports medicine (Auckland, NZ)*. 2018;48(11):2607–22.
- Lucassen PJ, Scheper W, Van Someren EJ. Adult neurogenesis and the unfolded protein response; new cellular and molecular avenues in sleep research. *Sleep medicine reviews*. 2009;13(3):183–6.
- McEwen BS. Protective and damaging effects of stress mediators: central role of the brain. *Dialogues Clin Neurosci*. 2006;8(4):367–81.
- Lucassen PJ, Meerlo P, Naylor AS, van Dam AM, Dayer AG, Fuchs E, et al. Regulation of adult neurogenesis by stress, sleep disruption, exercise and inflammation: Implications for depression and antidepressant action. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*. 2010;20(1):1–17.

38. Lou SJ, Liu JY, Chang H, Chen PJ. Hippocampal neurogenesis and gene expression depend on exercise intensity in juvenile rats. *Brain research*. 2008;1210:48–55.
39. Ernst C, Olson AK, Pinel JP, Lam RW, Christie BR. Antidepressant effects of exercise: evidence for an adult-neurogenesis hypothesis? *Journal of psychiatry & neuroscience : JPN*. 2006;31(2):84–92.
40. Craft LL, Perna FM. The Benefits of Exercise for the Clinically Depressed. *Prim Care Companion J Clin Psychiatry*. 2004;6(3):104–11.
41. Brand S, Colledge F, Ludyga S, Emmeneger R, Kalak N, Sadeghi Bahmani D, et al. Acute Bouts of Exercising Improved Mood, Rumination and Social Interaction in Inpatients With Mental Disorders. *Front Psychol*. 2018;9:249.
42. Ekkekakis P. The Dual-Mode Theory of affective responses to exercise in metatheoretical context: I. Initial impetus, basic postulates, and philosophical framework. *International Review of Sport and Exercise Psychology*. 2009;2(1):73–94.
43. Searle A, Calnan M, Lewis G, Campbell J, Taylor A, Turner K. Patients' views of physical activity as treatment for depression: a qualitative study. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 2011;61(585):149–56.
44. Dietrich A. Functional neuroanatomy of altered states of consciousness: The transient hypofrontality hypothesis. *Consciousness and Cognition*. 2003;12(2):231–56.
45. Dietrich A, Sparling PB. Endurance exercise selectively impairs prefrontal-dependent cognition. *Brain and Cognition*. 2004;55(3):516–24.
46. Ludyga S, Gerber M, Brand S, Holsboer-Trachsler E, Puhse U. Acute effects of moderate aerobic exercise on specific aspects of executive function in different age and fitness groups: A meta-analysis. *Psychophysiology*. 2016;53(11):1611–26.
47. Ludyga S, Gerber M, Puhse U, Looser VN, Kamijo K. Systematic review and meta-analysis investigating moderators of long-term effects of exercise on cognition in healthy individuals. *Nature human behaviour*. 2020.
48. Kubesch S, Bretschneider V, Freudemann R, Weidenhammer N, Lehmann M, Spitzer M, et al. Aerobic endurance exercise improves executive functions in depressed patients. *The Journal of clinical psychiatry*. 2003;64(9):1005–12.
49. Kramer AF, Colcombe S. Fitness Effects on the Cognitive Function of Older Adults: A Meta-Analytic Study-Revisited. *Perspectives on psychological science : a journal of the Association for Psychological Science*. 2018;13(2):213–7.
50. Bandura A. Self-efficacy: The exercise of control. New York, NY, US: W H Freeman/Times Books/ Henry Holt & Co; 1997. ix, 604–ix, p.
51. Stevens CJ, Bryan AD. Rebranding exercise: there's an app for that. *American journal of health promotion : AJHP*. 2012;27(2):69–70.
52. Carta MG, Hardoy MC, Pilu A, Sorba M, Floris AL, Mannu FA, et al. Improving physical quality of life with group physical activity in the adjunctive treatment of major depressive disorder. *Clinical practice and epidemiology in mental health : CP & EMH*. 2008;4:1.
53. Roshanaei-Moghaddam B, Katon WJ, Russo J. The longitudinal effects of depression on physical activity. *General Hospital Psychiatry*. 2009;31(4):306–15.
54. Schuch F, Vancampfort D, Firth J, Rosenbaum S, Ward P, Reichert T, et al. Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. *Journal of affective disorders*. 2017;210:139–50.
55. Schuch FB, Vancampfort D, Firth J, Rosenbaum S, Ward PB, Silva ES, et al. Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *The American journal of psychiatry*. 2018;175(7):631–48.
56. Bauman AE, Reis RS, Sallis JF, Wells JC, Loos RJ, Martin BW. Correlates of physical activity: why are some people physically active and others not? *Lancet (London, England)*. 2012;380(9838):258–71.
57. Vancampfort D, Stubbs B, Sienaert P, Wyckaert S, De Hert M, Rosenbaum S, et al. What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. *Psychiatria Danubina*. 2015;27(3):210–24.
58. Fuchs R, Göhner W, Seelig H. Effects of standardized group intervention on physical exercise and health: The MoVo-concept. *Journal of Physical Activity and Health*. 2011(8):794–803.
59. Fuchs R. Aufbau eines körperlich-aktiven Lebensstils im Kontext der medizinischen Rehabilitation: Ein motivational-volitionales Interventionskonzept (MoVo-LISA Projekt). Unveröffentlicher Endbericht Freiburg: Universität Freiburg. 2008.
60. Rosqvist E, Heikkinen E, Lyra TM, Hirvensalo M, Kallinen M, Leinonen R, et al. Factors affecting the increased risk of physical inactivity among older people with depressive symptoms. *Scandinavian journal of medicine & science in sports*. 2009;19(3):398–405.
61. Cohen-Mansfield J, Marx MS, Guralnik JM. Comparison of exercise models in an elderly population. *Aging clinical and experimental research*. 2006;18(4):312–9.
62. Krämer L, Helmes A, Bengel J. Understanding activity limitations in depression: Integrating the concepts of motivation and volition from health psychology into clinical psychology. *European Psychologist*. 2014(19):278–88.
63. Kramer LV, Helmes AW, Seelig H, Fuchs R, Bengel J. Correlates of reduced exercise behaviour in depression: the role of motivational and volitional deficits. *Psychol Health*. 2014;29(10):1206–25.
64. Firth J, Rosenbaum S, Stubbs B, Górczynski P, Yung AR, Vancampfort D. Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. *Psychological medicine*. 2016;46(14):2869–81.
65. Kerse N, Hayman KJ, Moyes SA, Peri K, Robinson E, Dowell A, et al. Home-based activity program for older people with depressive symptoms: DeLLITE--a randomized controlled trial. *Annals of family medicine*. 2010;8(3):214–23.
66. Mason OJ, Holt R. Mental health and physical activity interventions: a review of the qualitative literature. *Journal of mental health (Abingdon, England)*. 2012;21(3):274–84.
67. Fischer X, Kreppke JN, Zahner L, Gerber M, Faude O, Donath L. Telephone-Based Coaching and Prompting for Physical Activity: Short- and Long-Term Findings of a Randomized Controlled Trial (Movingcall). *International journal of environmental health and public health*. 2019;16(14).
68. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci*. 2011;6:42.
69. Gerber M, Beck J, Brand S, Cody R, Donath L, Eckert A, et al. The impact of lifestyle Physical Activity Counselling in IN-PATIENTS with major depressive disorders on physical activity, cardiorespiratory fitness, depression, and cardiovascular health risk markers: study protocol for a randomized controlled trial. *Trials*. 2019;20(1):367.